

- Four days prior to your procedure: Avoid eating seeds, nuts, corn, and any other form of roughage (lettuce, vegetables, bran cereals, etc.) or fiber. **DO NOT TAKE METAMUCIL or CITRUCEL.**
- Clear Liquids: It is important that large volumes of clear liquids be consumed the day before your procedure up to midnight to avoid dehydration. **Do not eat or drink after midnight before the procedure,** except for the liquid required for your bowel preparation. You may use a topical ointment such as Preparation H or Anusol to avoid irritation.
- **BREAKFAST DAY BEFORE PROCEDURE**
May include any of the following items:
 - One boiled or poached egg or small portion of skinless chicken, turkey, or fish.
 - White toast, **NO BUTTER**
 - One 8oz. can of Ensure, **Do not take Ensure Plus.**
 - Any clear liquid examples are:
 - Soft drinks*- Ginger Ale, Sprite, 7 Up, Gatorade, and Kool-Aid **NO RED OR PURPLE DYES.**
 - Strained fruit juice without pulp* – apple juice, white grape juice, lemonade
 - Water, Tea, or Coffee* -**NO MILK OR NON-DAIRY CREAMER,** sweeteners are permissible.
 - Soups*- low sodium chicken or beef bouillon/broth
 - Desserts*- hard candies Jell-O (lemon or lime) **NO FRUIT TOPPINGS,** Popsicles **NO SHERBETS OR FRUIT BARS.**
- **LUNCH DAY BEFORE PROCEDURE**
You may drink all of the clear liquids you would like. No solid food until after your Colonoscopy is performed.
- **DINNER DAY BEFORE PROCEDURE**
You may drink all of the clear liquids you would like. No solid food until after your Colonoscopy is performed.

SPECIAL – 7 DAY PREP INSTRUCTIONS FOR
COLONOSCOPY

7 DAYS BEFORE COLONOSCOPY

**2 TBSPS MILK OF MAGNESIA –CONTINUE TWICE A DAY
UNTIL 2 DAYS PRIOR TO YOUR COLONOSCOPY**

2 DAYS BEFORE COLONOSCOPY

1 BOTTLE MAGNESIUM CITRATE

DAY BEFORE COLONOSCOPY

MOVIPREP – AS DIRECTED

REMEMBER: STOP IRON 5 DAYS PRIOR TO PROCEDURE